How Diabetes Pills Work

Type II diabetes affect multiple organs in the body and diabetes pills work on specific organ(s) to lower blood glucose and HbA1c, which measures average level of blood sugar over 3 months. You will learn about different function, usage and benefit of the diabetes pills below. **Consult with your doctor about the best diabetes medication for you.**

Liver

Metformin (Glucophage)/ (Glucophage XR) Function: Lower the glucose release from liver to the blood Usage: Take medication with meals to reduce stomach upset Benefit: Lower A1c 1.0 - 2.0%

Liver and Pancreas

<u>Sitagliptin (Januvia), Saxagliptin</u> (Onglyza), Linagliptin (Tradjenta), Alogliptin (Nesina)

Function: Help extend the action of the gut hormones and slow down digestion

Usage: Take at the same time each day

Benefit: Lower A1c 0.6- 0.8%

Stomach

<u>Semaglutide (Rybelsus)</u>

Function: Making you feel full, lowering liver glucose output, preventing glucose formation in the liver and stimulating glucose uptake by cells Usage: Take it on empty stomach when you first wake up with a sip of plain water Benefits: lowering A1c 0.9-1.4%,

weight lost > 4 lb

Muscle Cell

<u>Pioglitazone (Actos)</u> **Function:** Require less insulin to process glucose or increase insulin sensitivity **Usage:** Take at the same time each day **Benefit:** Lower A1c 0.5 - 1.0%



To learn more about our doctor's group, call Member Relations: (415) 590-7418

Pancreas

Glimepiride (Amaryl), Glipizide (Glucotrol, Glucotrol XL), Glyburide (Micronase, Diabeta, Glynase), Nateglinide (Starlix), Repaglinide (Prandin) Function: Increase insulin release Usage: take medication before meals to reduce side effect Benefit: Lower A1c 1.0 - 2.0%

Kidneys

Canagliflozin (Invokana), Dapagliflozin (Farxiga), Empagliflozin (Jardiance) Function: Allow kidneys to urinate the excess sugars from the body Usage: Take medication before breakfast Benefit: Lower A1c 0.8 - 1.0%, lower weight 1-3 lbs

Intestines

Acarbose (Precose), Miglitol (Glyset) Function: Slow down the carbohydrate absorption from the intestine Usage: Take medication with 1st bite of the meal to improve the benefits of medication Benefit: Lower A1c 0.5 - 1.0%



f

aamgdoctors.com



facebook.com/DoctorsAAMG